

Contribution to the floristic knowledge of the Maddalena Mountains (Basilicata and Campania, southern Italy)

Leonardo Rosati¹, Vito Antonio Romano¹, Fabrizio Bartolucci², Liliana Bernardo³, Daniela Bouvet⁴, Laura Cancellieri⁵, Giuseppe Caruso⁶, Fabio Conti², Francesco Faraoni⁷, Enrico Banfi⁸, Gabriele Galasso⁸, Edda Lattanzi⁹, Paolo Lavezzi¹⁰, Simonetta Peccenini¹¹, Enrico Vito Perrino¹², Giovanni Salerno⁷, Adriano Sciandra¹³, Adriano Soldano¹⁴, Adriano Stinca^{15,16}, Chiara Totta⁷, Simonetta Fascetti³

1 School of Agricultural, Forestry, Food and Environmental Sciences, University of Basilicata, Potenza, Italy

2 Scuola di Bioscienze e Medicina Veterinaria, Università di Camerino/Centro Ricerche Floristiche dell'Appennino (Parco Nazionale del Gran Sasso e Monti della Laga), Italy **3** Department of Biology, Ecology and Earth Science, Università della Calabria, Rende, Italy **4** Department of Life Sciences and Systems Biology, Università di Torino, Torino, Italy **5** Department of Agriculture and Forestry Science, Tuscia University, Viterbo, Italy

6 Department of Agricultural, Food and Environmental Sciences, Università Politecnica delle Marche, Ancona, Italy **7** Department of Science, Università degli Studi Roma Tre, Rome, Italy **8** Museo di Storia Naturale di Milano, Sezione di Botanica, Milano, Italy **9** Via V. Cerulli 59, 00143 Roma, Italy **10** Via Teodosio Macrobio 19, 00136 Roma, Italy **11** DISTAV, Università di Genova, Genova, Italy **12** Mediterranean Agronomic Institute of Bari, Valenzano (Bari), Italy **13** Corso Carlo Brunet, 5A, Cuneo, Italy **14** Largo Brigata Cagliari 6, 13100 Vercelli, Italy **15** Department of Agriculture, University of Naples Federico II, Portici (Naples), Italy

16 Department of Environmental Biological and Pharmaceutical Sciences and Technologies, University of Campania Luigi Vanvitelli, Caserta, Italy

Corresponding author: Leonardo Rosati (leonardo.rosati@unibas.it)

Academic editor: L.Peruzzi | Received 3 March 2017 | Accepted 6 May 2017 | Published 31 May 2017

Citation: Rosati L, Romano VA, Bartolucci F, Bernardo L, Bouvet D, Cancellieri L, Caruso G, Conti F, Faraoni F, Banfi E, Galasso G, Lattanzi E, Lavezzi P, Peccenini S, Perrino EV, Salerno G, Sciandra A, Soldano A, Stinca A, Totta C, Fascetti S (2017) Contribution to the floristic knowledge of the Maddalena Mountains (Basilicata and Campania, southern Italy). Italian Botanist 3: 73–82. doi: 10.3897/italianbotanist.3.12519

Abstract

The inventory of the taxa collected during the annual field trip of the working group for Floristics, Systematics and Evolution of the Italian Botanical Society is reported. It was held in 2013 along the Maddalena Mountains, a mountain ridge of the southern Apennines located between the Basilicata and Campania administrative regions (southern Italy), considered as being poorly characterized in terms of vascular flora. A total of 701 units belonging to 74 plant families were recorded including two varieties and four hybrids.

Thirty-five taxa resulted endemic to Italy and only 11 alien species were detected, while 36 taxa are new or confirmed for the regional floras of Basilicata and/or Campania. In particular, 12 taxa are new for Basilicata, while four are confirmed. Regarding Campania, 14 taxa resulted new for the regional flora and five were confirmed.

Keywords

New floristic records, regional flora, southern Apennines, vascular flora

Introduction

In recent years, the working group for Floristics, Systematics and Evolution of the Italian Botanical Society has been very active in increasing the knowledge about the vascular flora of poorly known areas. Specifically, several contributions have been published regarding southern Italy (e.g. Bernardo et al. 2012, Wagensommer et al. 2014, Domina et al. 2015), as the result of the annual field trip organized by the group. Two of these contributions were focused on Basilicata (Conti et al. 2006, 2007b) and one on Campania region (Santangelo et al. 2010).

The selection of the areas to be investigated has been mostly addressed to fill the gaps pointed out by Scoppola and Blasi (2005) or by others regional or national projects, as the mapping of Important Plant Areas (Blasi et al. 2011).

In this paper, we present the results of the field trip held in 2013 in the southern Apennines and organized by the botanists of the University of Basilicata (L. Rosati, S. Fascetti and V.A. Romano), aimed at increasing our floristic knowledge of the western border between the Italian administrative regions of Basilicata and Campania.

Study area

The investigated area includes the central part of the Maddalena Mountains and some neighbouring biotopes of particular interest for vascular plants (Suppl. material 1: 1).

The Maddalena Mountains are a carbonatic ridge, aligned from NNW to SSE and extending for 40 km. They are located between the intramountain basins of the Val d'Agri (Basilicata) to the east and the Vallo di Diano (Campania) to the west. The altitude ranges from 300 m a.s.l. at the bottom of Melandro Valley, to 1503 m a.s.l. at the top of Serra Longa.

This sector of the southern Apennines is characterized by a remarkable geological complexity, mainly due to the effects of a translational tectonic that placed the formations of the Mesozoic carbonatic platform on the silico-clastic Lagonegrese Units. The Holocene detritus represents the connection with the recent alluvial deposits of the main valleys due to tectonic morpho-structures shaped like a “graben” (Grimaldi and Summa 2005).

The morpho-structure of the Maddalena Mountains is less affected by karst phenomena than other massifs of southern Italy; however, karstification is well developed in those areas where Cretaceous limestones prevail on Triassic dolomites (Celico 1979). Several tec-

tonic-karst basins (Mandrano, Mandranello, Magorno etc.) that are periodically flooded during the winter-spring season and are dotted with numerous sinkholes can be observed; they correspond to graben or contacts between lithotypes with different permeabilities.

Rainfall in the area is concentrated in the autumn-winter period, with a maximum in November-December and a minimum in summer (July-August); a summer drought of two to three months also occurs (Suppl. material 1: 2). Annual average temperature ranges from 12.5 to 14.7 °C, with the hottest months in summer (July-August) and the coldest in winter (January-February).

According to the classification proposed by Rivas-Martinez et al. (2011), a dominant Mediterranean pluviseasonal oceanic-semicontinental macrobioclimate can be recognized in the hilly and submontane belt (ranging from the lower humid mesomediterranean to humid supramediterranean phytoclimatic belt). However, local factors (increasing altitude, northern slopes) influence the shift towards the oceanic humid supratemperate belt.

The Maddalena Mountains are partially included within the Appennino Lucano National Park and within two Natura 2000 sites: "Monti della Maddalena" (code IT8050034) and "Faggeta di Moliterno" (code IT9210110). Nevertheless, this territory has never been the object of specific botanical contributions, except for the biotope "Faggeta di Moliterno" located in the southern part of the ridge (Fascetti et al. 2013). Other information concerning the vascular flora of this sector is available for the adjacent areas of the Lucanian Apennines (e.g. Gavioli 1948) or for the Cilento, Vallo di Diano and Alburni National Park (e.g. Moggi 2002, Rosati et al. 2010, 2012 and references therein).

Materials and methods

To maximize vascular flora sampling, 12 sites were selected as representative of the local diversity in terms of climate, litho-morphology, and land use (Suppl. material 1: 3). They were intensively sampled during the period 5–8 June, 2013 by 24 participants (Suppl. material 1: 4). The floristic list followed the same methodology as in previous contributions (e.g. Conti et al. 2006, Peccenini et al. 2010, Peruzzi et al. 2011).

The work was coordinated and the floristic list drawn up by the organizers with the contribution of all participants to the excursion. A revision of the samples collected during the field work was carried out at the University of Roma Tre (27–28 February, 2014), followed by specific studies and comparisons of unidentified taxa. Some critical samples were sent to specialists for determination: *Viola* (A. Scoppola, Viterbo), *Orobanche* (G. Domina, Palermo), ferns (D. Marchetti, Massa).

Nomenclature and taxa delimitation followed the updated version of the Checklist of Italian Flora (Bartolucci et al. 2016, Galasso et al. 2016), which is currently in the final stage of drafting, except for varieties (not considered in the Italian Checklist) and some hybrids (for further details see the notes in Suppl. material 1: 6). For the Orchidaceae, we followed Hertel and Presser (2015) and GIROS (2016), whereas for the genus *Rosa* we referred to Klastersky (1968).

For each unit at least one herbarium specimen was prepared and preserved in public or private herbaria listed in Suppl. material 1: 5.

The floristic list (Suppl. material 1: 6) was sorted in a linear sequence of families according to PPG I (2016), Christenhusz et al. (2011) and APG IV (2016) and taxa were ordered alphabetically. Synonyms are indicated in square brackets only in the case of changes that have occurred from the last version of the Italian Checklist of vascular flora (Conti et al. 2005, 2007a).

For each unit, we reported locations of collection, using the abbreviations given in Suppl. material 1: 3, and the prevailing environments of growth. The herbaria where samples are stored (acronyms for official herbaria according to Thiers 2016) are reported in brackets. The floristic novelties for the regional flora are marked with asterisks in Supplementary data (* = confirmed taxon, ** = taxon new for the regional flora, BAS = Basilicata; CAM = Campania). The letters “E” and “A” preceding the scientific names of the taxa indicate endemic and alien taxa, respectively.

Results

In total, more than 2600 samples of vascular plants were collected, belonging to 701 taxa and 74 families (see the complete floristic list in the Suppl. material 1: 1), including two varieties (*Ophrys apifera* Huds var. *bicolor* (Nägeli) E.Nelson and *Salvia officinalis* L. var. *angustifolia* Ten.) and four hybrids (*Acer × coriaceum* Bosc ex Tausch., *Anacamptis morio* (L.) R.M.Bateman, Pridgeon & M.W.Chase × *A. laxiflora* (Lam.) R.M.Bateman, Pridgeon & M.W.Chase, *Crataegus × media* Bechst. nothovar. *sicula* (K.Koch) K.I.Chr., and *Thymus longicaulis* C.Presl × *Thymus striatus* Vahl).

Thirty-five *taxa* are considered endemic to Italy (Peruzzi et al. 2014) and, amongst these, the following are restricted to southern Italy:

- *Achillea rupestris* Huter, Porta & Rigo subsp. *calcarea* (Huter, Porta & Rigo) Greuter
- *Alyssum diffusum* Ten. subsp. *calabicum* Španiel, Marhold, N.G.Passal. & Lihová
- *Epipactis collaris* S.Hertel
- *Epipactis lucana* H.Presser, S.Hertel & V.A.Romano
- *Knautia lucana* Lacaita & Szabó
- *Koeleria lucana* Brullo, Giusso & Miniss.
- *Koeleria splendens* C.Presl
- *Lathyrus jordanii* (Ten.) Ces., Pass. & Gibelli
- *Scorzonera villosa* Scop. subsp. *columnae* (Guss.) Nyman
- *Viola aethnensis* (DC.) Strobl subsp. *splendida* (W.Becker) Merxm. & Lippert

Only 11 taxa were alien (*Agrostemma githago* L., *Centaurea cyanus*, L., *Erigeron sumatrensis* Retz., *Gladiolus italicus* Mill., *Isatis tinctoria* L. subsp. *tinctoria*, *Malus pumila* Mill., *Papaver dubium* L. subsp. *dubium*, *Petroselinum crispum* (Mill.) Fuss, *Prunus dulcis* (Mill.) D.A.Webb, *Senecio inaequidens* DC., *Veronica persica* Poir.).

Thirty-six taxa have to be considered as floristic novelties because either new or confirmed for the regional flora of Basilicata and/or Campania.

In particular, 12 resulted new for the flora of Basilicata:

- *Asparagus tenuifolius* Lam.
- *Blackstonia acuminata* (W.D.J.Koch & Ziz) Domin subsp. *aestiva* (K.Malý) Zeltner
- *Carex humilis* Leyss.
- *Colchicum lusitanum* Brot.
- *Iberis umbellata* L.
- *Jasione montana* L.
- *Koeleria splendens* C.Presl
- *Lupinus albus* L. subsp. *graecus* (Boiss. & Spruner) Franco & P.Silva
- *Rosa inodora* Fr.
- *Rosa mollis* Sm.
- *Valerianella microcarpa* Loisel.
- *Viola eugeniae* Parl. subsp. *eugeniae*

As regards the following units, the subspecific rank for Basilicata was specified:

- *Rhinanthus alectorolophus* (Scop.) Pollich subsp. *alectorolophus*
- *Silene italica* (L.) Pers. subsp. *sicula* (Ucria) Jeanm.

Four taxa were confirmed for Basilicata:

- *Cardamine amporitana* Sennen & Pau
- *Thalictrum simplex* L. subsp. *simplex*
- *Thymus moesiacus* Velen.
- *Scabiosa columbaria* L. subsp. *portae* (Huter) Hayek

Fourteen taxa resulted new for the regional flora of Campania:

- *Bromus hordeaceus* L. subsp. *pseudothominei* (P.M.Sm.) H.Scholz
- *Carex tomentosa* L.
- *Hordeum geniculatum* All.
- *Juncus tenageia* L.f. subsp. *tenageia*
- *Knautia lucana* Lacaita & Szabó
- *Koeleria lucana* Brullo, Giusso & Miniss.
- *Pilosella piloselloides* (Vill.) Soják subsp. *praealta* (Gochnat) S.Bräut. & Greuter
- *Ranunculus peltatus* Schrank subsp. *peltatus*
- *Rosa mollis* Sm.
- *Rubus incanescens* L.
- *Sanguisorba officinalis* L.
- *Scabiosa columbaria* L. subsp. *portae* (Huter) Hayek
- *Silene italica* (L.) Pers. subsp. *sicula* (Ucria) Jeanm.
- *Trifolium phleoides* Willd.

Finally, five taxa were confirmed for Campania:

- *Alisma lanceolatum* With.
- *Alopecurus aequalis* Sobol.
- *Myosotis nemorosa* Besser
- *Sabulina glaucina* (Dvořáková) Dillenb. & Kadereit
- *Vicia serratifolia* Jacq.

Conclusions

The high number of taxa surveyed in a few days and in a limited number of sampling localities undoubtedly indicates the high level of biodiversity of the Maddalena Mountains, an area that until now did not attract explorations by botanists.

The number of new or confirmed units at the regional level underlines the fact that floristic knowledge of Basilicata and Campania cannot yet be considered satisfactory, despite numerous publications produced in recent years (e.g. Azzella et al. 2014, Bernardo and Caldararo 2014, Rosati et al. 2012, 2015, Bonari et al. 2016, Roma-Marzio et al. 2016, Stinca et al. 2016, 2017). It should be emphasized that several collected units are linked to wetlands that risk to disappear or to become altered both globally and locally. Finally, the limited number of surveyed exotic species can be considered as an indicator of the favourable conservation status of the investigated territories that are characterized mostly by natural/semi-natural habitats or by traditional arable land of mountainous areas.

Acknowledgements

We are grateful to Anna Scoppola, Dino Marchetti, Giannantonio Domina for the determination of some critical samples. Giovanna Potenza and Emmanuele Farris kindly collaborate to organize the field trip. A special thanks to Michele Cicerchia and his family for the warm hospitality at the Hotel Eden in Brienza. The visit to the hill of Caracciolo's Castle was a courtesy of the Municipality of Brienza. We should also like to thank Fernando Lucchese and Alfred Mayer for having hosted and supported the working days for review of the critical samples at the University of Roma Tre. Michela Marignani has kindly revised the English language. We are grateful to Francesco Roma-Marzio and Giannantonio Domina for the improvements to the first draft of the paper.

References

- APG IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society* 181: 1–20. <https://doi.org/10.1111/boj.12385>

- Azzella MM, Rosati L, Iberite M, Bolpagni R, Blasi C (2014) Changes in aquatic plants in the Italian volcanic-lake system detected using current data and historical records. *Aquatic Botany* 112: 41–47. <https://doi.org/10.1016/j.aquabot.2013.07.005>
- Bartolucci F, Peruzzi L (2014) *Thymus paronychoides* (Lamiaceae), a neglected species from Sicily belonging to *Thymus* sect. *Hyphodromi*. *Folia Geobotanica* 49: 83–106. <https://doi.org/10.1007/s12224-013-9156-2>
- Bartolucci F, Walter J (2015) Typification of names in the genus *Thymus* (Lamiaceae). *Phytotaxa* 221(2): 137–147. <https://doi.org/10.11646/phytotaxa.221.2.3>
- Bartolucci F, Peruzzi L, Passalacqua N (2013) Taxonomic notes and typification of names within the genus *Thymus* L. (Lamiaceae). *Taxon* 62(6): 1308–1314. <https://doi.org/10.12705/626.1>
- Bartolucci F, Peruzzi L, Galasso G, Conti F (Eds) 2016) Checklist aggiornata della flora vascolare autoctona d'Italia. *Notiziario della Società Botanica Italiana* 0: 29–30.
- Bernardo L, Caldararo F (2014) Nuove acquisizioni per la flora della Basilicata. *Informatore Botanico Italiano* 46(1): 53–57.
- Bernardo L, Bartolucci F, Cancellieri L, Costalonga S, Galasso G, Galesi R, Gargano D, Iberite M, Iocchi M, Lattanzi E, Lavezzo P, Magrini S, Peccenini S, Sciandrello S, Scoppola A, Signorino G, Tilia A, Spampinato G (2012) Contributo alla conoscenza floristica della Calabria: resoconto dell'escursione del Gruppo di Floristica (S.B.I.) nel 2008 nella Presila catanzarese. *Informatore Botanico Italiano* 44(1): 125–151.
- Blasi C, Marignani M, Copiz R, Fipaldini M, Bonacquisti S, Del Vico E, Rosati L, Zavattero L (2011) Important Plant Areas in Italy: From data to mapping. *Biological Conservation* 144: 220–226. <https://doi.org/10.1016/j.biocon.2010.08.019>
- Bonari G, Cambria S, Rosati L, Domina G (2016) Contributo alla conoscenza della distribuzione di *Monotropa hypophaea* Wallr. (Ericaceae) in Italia. *Atti della Società Toscana di Scienze Naturali, Memorie Serie B* 122(2015): 11–16.
- Castroviejo S (Ed.) 2015. Flora iberica 9. Real Jardín Botánico, CSIC, Madrid.
- Celico P (1979) Considerazioni sull'idrogeologia di alcune zone dell'Italia centro-meridionale alla luce dei risultati di recenti indagini geognostiche. *Memorie e Note Istituto Geologia Applicata* 15: 1–43.
- Christensen KI (1992) Revision of *Crataegus* Sect. *Crataegus* and Nothosect. *Crataeguineae* (Rosaceae-Maloideae) in the Old World. *Systematic Botany Monographs* 35: 1–199. <https://doi.org/10.2307/25027810>
- Christenhusz MJM, Reveal JL, Farjon A, Gardner MF, Mill RR, Chase MW (2011) A new classification and linear sequence of extant gymnosperms. *Phytotaxa* 19: 55–70. <https://doi.org/10.11646/phytotaxa.19.1.3>
- Conti F, Abbate G, Alessandrini A, Blasi C [Eds] (2005) An annotated checklist of the Italian vascular flora. Palombi Editori, Roma, 420 pp.
- Conti F, Angiolini C, Bernardo L, Costalonga S, Di Pietro R, Fascetti S, Giardina G, Giovi E, Gubellini L, Lattanzi E, Lavezzo P, Peccenini S, Salerno G, Scoppola A, Tinti D, Turrisi R (2006) Contributo alla conoscenza floristica della Basilicata: resoconto dell'escursione del gruppo di floristica (S.B.I.) nel 2003. *Informatore Botanico Italiano* 38(2): 383–409.
- Conti F, Alessandrini A, Bacchetta G, Banfi E, Barberis G, Bartolucci F, Bernardo L, Bonacquisti S, Bouvet D, Bovio M, Brusa G, Del Guacchio E, Foggi B, Frattini S, Galasso G, Gallo

- L, Gangale C, Gottschlich G, Grünanger P, Gubellini L, Iriti G, Lucarini D, Marchetti D, Moraldo B, Peruzzi L, Poldini L, Prosser F, Raffaelli M, Santangelo A, Scassellati E, Scortegagna S, Selvi F, Soldano A, Tinti D, Ubaldi D, Uzunov D, Vidali M (2007a) Integrazioni alla checklist della flora vascolare italiana. *Natura Vicentina* 10: 5–74.
- Conti F, Bartolucci F, Tinti D, Bernardo L, Costalunga S, Lattanzi E, Lavezzo P, Salerno G, Fasceri S, Iocchi M, Mele C, Tardella FM (2007b) Secondo contributo alla conoscenza floristica della Basilicata: resoconto dell'escursione del Gruppo di Floristica (S.B.I.) nel 2004. *Informatore Botanico Italiano* 31(1): 11–33.
- Domina G, Marino P, Castellano G, Amato F, Cambria S, Cancellieri L, Crisafulli A, Cristaudo A, Faraoni F, Galesi R, Guarino R, Lattanzi E, Lavezzo P, Longo D, Maiorca G, Peccenini S, Perrino E, Salerno G, Scolastri A, Soldano A, Stinca A, Wagensommer R, Xibilia L, Raimondo F (2015) Contributo alla conoscenza floristica dei monti Sicani (Sicilia): resoconto dell'escursione del Gruppo di Floristica (S.B.I.) nel 2012. *Informatore Botanico Italiano* 47(2): 155–177.
- Fasceri S, Pirone G, Rosati L (2013) The vegetation of the Maddalena Mountains (Southern Italy) *Plant Sociology* 50(2): 5–37.
- Galasso G, Bartolucci F, Peruzzi L, Ardenghi NMG, Banfi E, Celesti-Grapow L, Conti F (Eds) (2016) Checklist aggiornata della flora vascolare alloctona d'Italia. *Notiziario della Società Botanica Italiana* 0: 49–50.
- Gavioli O (1948) Synopsis Flora Lucanae. *Nuovo Giornale Botanico Italiano*, nuova serie 54: 134–135.
- GIROS (2016) Orchidee d'Italia. Il Castello. Milano.
- Grimaldi S, Summa G (2005) Caratteri idrogeologici ed idrogeochimici del settore meridionale dei Monti della Maddalena (Appennino Meridionale). *Giornale Geologia Applicata* 2: 348–356.
- Hertel S, Presser H (2015) Zur Kenntnis einiger *Epipactis*-Arten in Italien. *Berichte aus den Arbeitskreisen Heimische Orchideen*. 32(2): 79–109.
- Klastersky I (1968) *Rosa* L. In: Tutin TG, Burges NA, Chater AO, Edmondson JR, Heywood VH, Moore DM, Valentine DH, Walters SM, Webb DA (Eds) *Flora Europaea*. 2. Cambridge University Press.
- Moggi G (2002) Catalogo della Flora del Cilento (Salerno). Repertorio delle piante vascolari finora segnalate e problemi sistematici connessi. *Informatore Botanico Italiano* 33(Suppl. 3): 1–116.
- Peccenini S, Polatschek A (2016) The genus *Erysimum* (Brassicaceae) in Italy, part III: Key to the species and description of the new species *E. apenninum*, *E. etruscum*, *E. pignattii*. *Annalen des Naturhistorischen Museums in Wien. Serie B, Botanik und Zoologie* 118: 147–166.
- Peccenini S, Barberis G, Bartolucci F, Cancellieri L, Conti F, Costalunga S, Dente F, Iocchi M, Lattanzi E, Lavezzo P, Lupino F, Magrini S, Salerno G, Tardella FM, Terzo V, Tinti D, Zappa E (2010) Contributo alla conoscenza floristica della Liguria: resoconto dell'escursione del Gruppo di Floristica nel 2006 sulle Alpi Liguri meridionali. *Informatore Botanico Italiano* 42(1): 3–25.
- Peruzzi L, Barbo M, Bartolucci F, Bovio, M, Carta A, Ciccarelli D, Conti F, Costalunga S, Di Pietro R, Galasso G, Gestri G, Lattanzi E, Lavezzo P, Marsili S, Peccenini S, Pierini B, Tardella FM, Terzo V, Turrisi RE, Bedini G (2011) Contributo alla conoscenza floristica

- delle Colline Pisane: resoconto dell'escursione del Gruppo di Floristica (S.B.I.) nel 2009. Informatore Botanico Italiano 43(1): 3–27.
- Peruzzi L, Conti F, Bartolucci F (2014) An inventory of vascular plants endemic to Italy. Phytotaxa 168(1): 1–75. <https://doi.org/10.11646/phytotaxa.168.1.1>
- Peruzzi L, Domina G, Bartolucci F, Galasso G, Peccenini S, Raimondo FM, Albano A, Alessandrini A, Banfi E, Barberis G, Bernardo L, Bovio M, Brullo S, Brundu G, Brunu A, Camarda I, Carta L, Conti F, Croce A, Iamonic D, Iberite M, Iiriti G, Longo D, Marsili S, Medagli P, Pistarino A, Salmeri C, Santangelo A, Scassellati E, Selvi F, Soldano A, Stinca A, Villani MC, Wagensommer RP, Passalacqua NG (2015) An inventory of the names of vascular plants endemic to Italy, their loci classici and types. Phytotaxa 196: 1–217. <https://doi.org/10.11646/phytotaxa.196.1.1>
- Pignatti S (1982) Flora d'Italia 1–3. Edagricole, Bologna.
- PPG I (2016) A community-derived classification for extant lycophytes and ferns. Journal of Systematics and Evolution 54(6): 563–603. <https://doi.org/10.1111/jse.12229>
- Rivas Martínez S, Rivas Saenz S, Penas Merino A (2011) Worldwide Bioclimatic classification system. Global Geobotany 1: 1–638.
- Roma-Marzio F, Bernardo L, Liguori P, Peruzzi L (2016) Vascular flora of Monte Sparviere (Southern Italy, Pollino massif). Atti della Società Toscana di Scienze Naturali, Memorie serie B 122 (2015): 73–88.
- Rosati L, Filibeck G, De Lorenzis A, Lattanzi E, Surbera S, Fascetti S, Blasi C (2010) La vegetazione Forestale dei Monti Alburni, Parco Nazionale del Cilento e Vallo di Diano: analisi fitosociologica e significato fitogeografico. Fitosociologia 47(2): 17–55.
- Rosati L, Salerno G, Del Vico E, Lapenna MR, Villani M, Filesi L, Fascetti S, Lattanzi E (2012) Un aggiornamento della flora del Cilento e della Campania. Informatore Botanico Italiano 44(1): 111–119.
- Rosati L, Farris E, Tilia A, Potenza G, Fascetti S (2014) *Gymnospermium scipetarum* (Berberidaceae) specie nuova per la flora italiana. In Peruzzi L, Domina G (2014) Atti della Riunione scientifica del Gruppo di Floristica, Tassonomia ed Evoluzione della Società Botanica Italiana “Floristica, Sistematica ed Evoluzione”. Orto Botanico di Roma, La Sapienza, Università di Roma, 21–22 novembre 2014: 7–8.
- Santangelo A, Bernardo L, Bertani G, Bronzo E, Cancellieri L, Costalonga S, Croce A, Del Vico E, Fascetti S, Fortini P, Gangale C, Gubellini L, Iocchi M, Lapenna MR, Lattanzi E, Lavezzo P, Lupino F, Magrini S, Marino R, Paura B, Peccenini S, Peruzzi L, Rosati L, Salerno G, Scoppola A, Strumia S, Tardella FM, Uzunov D (2010) Contributo alla conoscenza floristica del Massiccio del Matese: resoconto dell'escursione del Gruppo di Floristica (S.B.I.) nel 2007. Informatore Botanico Italiano 42(1): 109–143.
- Scoppola A, Blasi C (Eds) (2005) Stato delle conoscenze sulla Flora Vascolare d'Italia. Palombi Editori, Roma.
- Stinca A, Croce A, D'Auria G, Salerno G, Santangelo A, Rosati L, Motti R (2016) Nuovi dati sulla flora vascolare aliena della Campania (Sud Italia). Atti della Società Toscana di Scienze Naturali, Memorie Serie B, 122(2015): 89–110.
- Stinca A, Ravo M, Giacanelli V, Conti F (2017) Integrazioni alla flora vascolare dell'Isola di Capri (Campania, Sud Italia). Atti della Società Toscana di Scienze Naturali, Memorie Serie B, 123(2016): 65–70.

- Thiers B (2016) Index Herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/> [accessed 15.1.2017]
- Wagensommer RP, Marrese M, Perrino EV, Bartolucci F, Cancellieri L, Carruggio F, Conti F, Di Pietro R, Fortini P, Galasso G, Lattanzi E, Lavezzo P, Longo D, Peccenini S, Rosati L, Russo G, Salerno G, Scoppola A, Soldano A, Stinca A, Tilia A, Turco A, Medagli P, Forte L (2014) Contributo alla conoscenza floristica della Puglia: resoconto dell'escursione del Gruppo di Floristica (S.B.I.) nel 2011 nel settore meridionale dei Monti della Daunia. Informatore Botanico Italiano 46(2): 175–208.

Supplementary material I

Supplementary data

Authors: Leonardo Rosati, Vito Antonio Romano, Fabrizio Bartolucci, Liliana Bernardo, Daniela Bouvet, Laura Cancellieri, Giuseppe Caruso, Fabio Conti, Francesco Faraoni, Gabriele Galasso, Edda Lattanzi, Paolo Lavezzo, Simonetta Peccenini, Enrico Vito Perrino, Giovanni Salerno, Adriano Sciandra, Adriano Soldano, Adriano Stinca, Chiara Totta, Simonetta Fascetti

Data type: Word .doc file

Explanation note:

1. Study area. Surveyed sites are marked with red squares (for details see Suppl. material 1: 3 and 7–8).
2. Thermo-pluviometric diagram of two representative stations of Maddalena Mountains. Observation period of Moliterno station: 1923–2007; Sala Consilina: 1975–1995. Data from “Annali idrologici, Ministero dei lavori pubblici, Servizio idrografico”. Solid line: average monthly values; dashed lined: average monthly maximum temperature; point-dashed line: average monthly minimum temperature.
3. Coded locality and geographical features of surveys. Coordinates are reported with decimal degrees WGS84. For detailed topographic map of each sites see Suppl. material 1: 7–8.
4. The 24 participants of the excursion of the Italian Botanical Society to the Maddalena Mountains, plus P. Scelzo, the Mayor of Brienza, (the second at the bottom from the left) during the visit to the mediaeval Caracciolo's Castle (courtesy of the Municipality of Brienza). Photo V.A. Romano.
5. Herbarium acronyms and institutions where the collected samples are stored.
6. Floristic list of taxa surveyed in the Maddalena Mountains.
7. Topographic map of surveyed sites (1–6). Floristic sampling sites are coded according to Supplementary 1 and 3; the cells grid is 1 × 1 km.
8. Topographic map of surveyed sites (7–12). Floristic sampling sites are coded according to Suppl. material 1: 1 and 3; the cells grid is 1 × 1 km.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.