

## Notulae to the Italian alien vascular flora: 5

Gabriele Galasso<sup>1</sup>, Giannantonio Domina<sup>2</sup>, Michele Adorni<sup>3</sup>,  
Nicola M.G. Ardenghi<sup>4</sup>, Gianmaria Bonari<sup>5</sup>, Sergio Buono<sup>6</sup>, Laura Cancellieri<sup>7</sup>,  
Giuseppina Chianese<sup>8</sup>, Giulio Ferretti<sup>9</sup>, Tiberio Fiaschi<sup>10</sup>, Luigi Forte<sup>11</sup>,  
Riccardo Guarino<sup>12</sup>, Rocco Labadessa<sup>13</sup>, Lorenzo Lastrucci<sup>9</sup>, Lorenzo Lazzaro<sup>9</sup>,  
Sara Magrini<sup>14</sup>, Luigi Minuto<sup>15</sup>, Sara Mossini<sup>16</sup>, Nicola Olivieri<sup>17</sup>, Anna Scoppola<sup>7</sup>,  
Adriano Stinca<sup>18</sup>, Claudia Turcato<sup>15</sup>, Chiara Nepi<sup>19</sup>

**1** Sezione di Botanica, Museo di Storia Naturale di Milano, Corso Venezia 55, 20121 Milano, Italy **2** Dipartimento di Scienze Agrarie, Alimentari e Forestali (SAAF), Università di Palermo, Viale delle Scienze, ed. 4, 90128 Palermo, Italy **3** Via degli Alpini 7, 43037 Lesignano de' Bagni (Parma), Italy **4** Dipartimento di Scienze della Terra e dell'Ambiente, Università di Pavia, Via Sant'Epifanio 14, 27100 Pavia, Italy **5** Department of Botany and Zoology, Masaryk University, Kotlarska 2, CZ-611 37 Brno, Czech Republic **6** Via XXV Aprile 6, 01010 Oriolo Romano (Viterbo), Italy **7** Dipartimento di Scienze Agrarie e Forestali (DAFNE), Università della Tuscia, Via San Camillo de Lellis snc, 01100 Viterbo, Italy **8** Musei delle Scienze Agrarie, Università di Napoli Federico II, Via Università 100, 80055 Portici (Napoli), Italy **9** Dipartimento di Biologia, Università di Firenze, Via G. La Pira 4, 50121 Firenze, Italy **10** Dipartimento di Scienze della Vita, Università di Siena, Via P.A. Mattioli 4, 53100 Siena, Italy **11** Dipartimento di Biologia e Museo Orto Botanico, Università di Bari Aldo Moro, Via E. Orabona 4, 70125 Bari, Italy **12** Sezione di Botanica ed Ecologia Vegetale, Dipartimento di Scienze e Tecnologie Biologiche, Chimiche e Farmaceutiche (STEBICEF), Università di Palermo, Via Archirafi 20, 90123 Palermo, Italy **13** Associazione Centro Studi de Romita, Via G. Postiglione 9, 70126 Bari, Italy **14** Banca del Germoplasma della Tuscia, Università della Tuscia, Largo dell'Università snc, blocco c, 01100 Viterbo, Italy **15** Dipartimento di Scienze della Terra, dell'Ambiente e della Vita (DISTAV), Università di Genova, Corso Dogali 1/m, 16136 Genova, Italy **16** Via E. Rovati 6, 27049 Stradella (Pavia), Italy **17** Via Maestri del Lavoro 40, 64100 Teramo, Italy **18** Dipartimento di Scienze e Tecnologie Ambientali, Biologiche e Farmaceutiche, Università della Campania Luigi Vanvitelli, Via A. Vivaldi 43, 81100 Caserta, Italy **19** Sezione di Botanica Filippo Parlatore, Museo di Storia Naturale, Università di Firenze, Via G. La Pira 4, 50121 Firenze, Italy

Corresponding author: Gabriele Galasso (gabriele.galasso@comune.milano.it)

---

Academic editor: Lorenzo Peruzzi | Received 17 April 2018 | Accepted 27 April 2018 | Published 14 May 2018

**Citation:** Galasso G, Domina G, Adorni M, Ardenghi NMG, Bonari G, Buono S, Cancellieri L, Chianese G, Ferretti G, Fiaschi T, Forte L, Guarino R, Labadessa R, Lastrucci L, Lazzaro L, Magrini S, Minuto L, Mossini S, Olivieri N, Scoppola A, Stinca A, Turcato C, Nepi C (2018) Notulae to the Italian alien vascular flora: 5. Italian Botanist 5: 45–56. <https://doi.org/10.3897/italianbotanist.5.25910>

---

## Abstract

In this contribution, new data concerning the distribution of vascular flora alien to Italy are presented. It includes new records, and confirmations for Italy or for Italian administrative regions of taxa in the genera *Albizia*, *Anredera*, *Bougainvillea*, *Cardamine*, *Cenchrus*, *Cephalaria*, *Ceratochloa*, *Cytisus*, *Datura*, *Delosperma*, *Euonymus*, *Freesia*, *Hylotelephium*, *Lantana*, *Musa*, *Physalis*, *Rotala*, *Styphnolobium*, *Trachycarpus*, and *Tradescantia*. Nomenclature and distribution updates, published elsewhere, and corrections are provided as supplementary material.

## Keywords

Alien species, floristic data, Italy, nomenclature

## How to contribute

The text for the new records should be submitted electronically to Chiara Nepi (chiara.nepi@unifi.it). The corresponding specimen along with its scan or photograph has to be sent to FI Herbarium: Sezione di Botanica Filippo Parlatore del Museo di Storia Naturale, Via G. La Pira 4, 50121 Firenze (Italy). Those texts concerning nomenclatural novelties (typifications only for accepted names), status changes, exclusions, and confirmations should be submitted electronically to: Gabriele Galasso (gabriele.galasso@comune.milano.it). Each text should be within 2,000 characters (spaces included).

## Floristic records

### *Albizia julibrissin* Durazz. (Fabaceae)

+ (CAS) **CAL**: Montegiordano (Cosenza), fraz. Montegiordano Marina, massicciata ferroviaria presso Via Canale G. Garibaldi (WGS84: 40.031813°N; 16.600047°E), massicciata ferroviaria, ca. 10 m, 21 August 2017, N. Olivieri (FI). – Casual alien species new for the flora of Calabria.

A young specimen of the species grows on pebbly ground at the railway roadbed crossing the village's suburban area near an overpass. It originated from the seeds produced by some trees cultivated along a neighboring road. The native range of *Albizia julibrissin* includes southern Caucasus, Iran, Pakistan, India, Nepal, Bangladesh, Sri Lanka, Myanmar, China, Korea, Japan, Ethiopia, Eritrea, and Somalia. The species was introduced in Italy for ornamental purposes in 1745 (Targioni Tozzetti 1896) and it is currently widely planted as ornamental in parks and gardens around the world.

N. Olivieri

*Anredera cordifolia* (Ten.) Steenis (Basellaceae)

+ (CAS) **MAR**: Cupra Marittima (Ascoli Piceno), presso la stazione ferroviaria (WGS84: 43.023750°N; 13.860402°E), incolto, ca. 6 m, 3 August 2017, *N. Olivieri* (FI). – Casual alien species confirmed for the flora of Marche.

At Cupra Marittima, this species grows on sandy soil at the edge of a small neglected area, along with *Broussonetia papyrifera* (L.) Vent. and *Ailanthus altissima* (Mill.) Swingle. The species was no longer recorded in the Marche after 1950 (Galasso et al. 2018). According to Viegi et al. (2004), reporting Brilli-Cattarini's personal communications, *Anredera cordifolia* was previously observed in Ancona in 1945, and in Pesaro in 1939.

N. Olivieri

*Bougainvillea spectabilis* Willd. (Nyctaginaceae)

+ (CAS) **ABR**: Pescara (Pescara), presso Viale V. Pepe (WGS84: 42.458347°N; 14.231300°E), bordo di marciapiede, ca. 3 m, NE, 9 August 2017, *N. Olivieri* (FI). – Casual alien species new for the flora of Abruzzo.

A young plant has developed on the inner edge of a sidewalk at the base of the perimetral wall of a private home in the urban area of Pescara. The location is partially shaded and at a short distance from the Adriatic Sea, but it is sheltered from sea winds by the buildings. In Italy, *Bougainvillea spectabilis* is also reported as a casual alien in Sardegna (Bacchetta et al. 2009, Galasso et al. 2018).

N. Olivieri

*Cardamine occulta* Hornem. (Brassicaceae)

+ (CAS) **LAZ**: Roma (Roma), lungo Via del Biscione (WGS84: 41.895524°N; 12.472969°E), aiuola, 18 m, 28 October 2017, *A. Stinca* (FI, PORUN). – Casual alien species new for the flora of Lazio.

In Italy, *Cardamine occulta* is recorded for Piemonte, Lombardia, Trentino-Alto Adige, Veneto, Friuli Venezia Giulia, Emilia-Romagna, Toscana, Campania, and Sardegna (Galasso et al. 2018). According to Stinca et al. (2017), this species has been largely spread by nurseries and soil transportation.

A. Stinca, G. Chianese

*Cenchrus setaceus* (Forssk.) Morrone (Poaceae)

+ (CAS) **TOS**: Piombino (Livorno), fraz. Baratti (WGS84: 42.998797°N; 10.516870°E), margine stradale, 8 m, 21 October 2017, *T. Fiaschi* (FI). – Casual alien species new for the flora of Toscana.

The occurrence of this species in Toscana does not seem to be linked to direct human cultivation. Seeds of *Cenchrus setaceus* are known to have a very efficient wind dispersal strategy. However, given that the nearest known populations occur a few hundred kilometres away from Baratti, wind is not the most likely vector. The species is included in the list of Invasive Alien Species of Union Concern (Commission Implementing Regulation EU 2016/1141) and was found in three separate nuclei 10 m apart, in a parking area of a renowned tourist location, where seeds could have been introduced *via* pet fur or car mats. According to Galasso et al. (2018), it occurs in Lazio, Puglia, Calabria, Sicilia, and Sardegna.

G. Bonari, T. Fiaschi, R. Guarino

#### *Cephalaria syriaca* (L.) Schrad. (Dipsacaceae)

+ (NAT) **LAZ**: Roma (Roma), Scalo ferroviario Ostiense (WGS84: 41.871550°N; 12.488699°E ± 700 m), ca. 20 m, 28 May 1954, A. Cacciato (RO); Tarquinia (Viterbo), loc. Pian di Spille, zona militare (WGS84: 42.255734°N; 11.678989°E), incolto erboso al margine di campo coltivato a fieno greco, 4 m, 17 April 2017, A. Scoppola (FI, UTV no. 35200); Monte Romano (Viterbo), valle del Fiume Mignone (WGS84: 42.251059°N; 11.877385°E), margine di campi lungo strada sterrata in luogo arido, 109 m, 20 April 2017, A. Scoppola, L. Cancellieri (UTV no. 35202, Herb. L. Cancellieri). – Naturalized alien species confirmed for the flora of Lazio.

*Cephalaria syriaca* belongs to the Mediterranean-Turanian element, growing as a weed in cereal fields and waste places (Matthews 1972). It is not listed for Lazio by Anzalone et al. (2010), while Lucchese (2017) records the species for this administrative region, but without providing information on recent records or herbarium specimens. Accordingly, it has been excluded from the confirmed regional alien flora (Galasso et al. 2018). It is not reported in the flora of the Pian di Spille coast (Iocchi and Bartolucci 2008). However, it was reported by Cacciato (1955) in the Ostiense railway station in Roma, an area deeply transformed due to the urban development of the last decades. This ancient discovery is attested by a 1954 herbarium specimen preserved in RO. However, this record was ignored by Lucchese (2017). In the two new localities, *C. syriaca* occurs with a conspicuous number of fertile individuals, spread at the edge of cultivated fields and waste places.

L. Cancellieri, A. Scoppola

#### *Ceratochloa cathartica* (Vahl) Herter (Poaceae)

+ (NAT) **PUG**: Bari (Bari), foce del Torrente Lamasinata (WGS84: 41.134654°N; 16.827638°E), prateria umida su substrato sabbioso, 1 m, 30 September 2017, R. Labadessa (FI); Bari (Bari), Torrente Lamasinata (WGS84: 41.134647°N;

16.827632°E), prati umidi su sabbia, 1 m, 30 September 2017, *R. Labadessa* (BI no. 40486, no. 40487). – Naturalized alien species new for the flora of Puglia.

*Ceratochloa cathartica* is an alien plant from South America, whose European distribution ranges from Portugal to Ukraine, and from Great Britain to Italy (Ryves et al. 1996). This species was probably introduced as fodder in Europe, where it may be found as naturalized in the warmer regions (Ryves et al. 1996). Several individuals were found within an area of about two hectares, in a wet sandy meadow dominated by annual and perennial grass species.

R. Labadessa, L. Forte

### *Cytisus striatus* (Hill) Rothm. (Fabaceae)

+ (NAT) **ITALIA (LIG):** Noli (Savona), fraz. Tosse, fra la SP8 e l'Autostrada dei Fiori A10 (WGS84: 44.227616°N; 8.393363°E), macchia mediterranea, 177 m, 20 June 2017, leg. *L. Minuto*, det. *C. Turcato* (FI, GE). – Naturalized alien species new for the flora of Italy (Liguria).

*Cytisus striatus* is native to Morocco, Portugal, and Spain. It has been introduced into a number of northwestern European countries (England, Scotland, Wales, and France), and in the Americas (California and Oregon in the U.S.A., and Argentina), where it is considered to be an invasive shrub. A large population, identified following the key published by Frodin and Heywood (1968), was detected growing on the motorway embankment. Other plants were observed along the Autostrada A10 from Albenga to Savona Vado, and along the Autostrada A12 near Sestri Levante (loc. Rocche di Sant'Anna).

C. Turcato, L. Minuto

### *Datura wrightii* Regel (Solanaceae)

+ (CAS) **LAZ:** Bracciano (Roma), fraz. Vigna di Valle, Museo Storico dell'Aeronautica Militare, presso l'Aeroporto di Vigna di Valle (WGS84: 42.085366°N; 12.218396°E), suolo sabbioso presso il lago, 158 m, 16 July 2017, *S. Buono* (FI, UTV). – Casual alien species new for the flora of Lazio.

*Datura wrightii* is a perennial plant native to the southwestern United States and Mexico. In southern Europe, it was widely confused with the closely related species *D. inoxia* Mill. (Verlooove 2008). Reports of the occurrence of this species as casual or naturalized alien in some Italian regions are very recent (Banfi and Galasso 2010, Verlooove et al. 2010, Ardenghi et al. 2011, Del Guacchio 2011, Cerutti and Motta 2012, D'Aleo and Bonanno 2016, Galasso et al. 2018). A single individual was observed growing on sandy soil near Lake Bracciano, along with other alien species as *Abutilon theophrasti* Medik., *Ludwigia peploides* (Kunth) P.H.Raven subsp. *montevidensis*

(Spreng.) P.H.Raven, *Pavonia hastata* Cav. (see also Galasso et al. 2017), and *Physalis peruviana* L. (see beyond in this contribution). The plant regularly developed a high number of flowers and fruits from July to December 2017.

S. Buono, S. Magrini, A. Scoppola

*Delosperma cooperi* (Hook.f.) L.Bolus (Aizoaceae)

+ (CAS) **EMR**: Castel San Giovanni (Piacenza), Via Fratelli Bandiera (SP10R), angolo con Via Bottarone (WGS84: 45.06085°N; 9.43153°E), ciglio stradale, con *Setaria italica* subsp. *viridis* e *Lactuca sativa* subsp. *serriola*, 75 m, 3 September 2017, N. Ardenghi, S. Mossini (FI). – Casual alien species new for the flora of Emilia-Romagna.

A single flowering individual was found in the growing site.

N.M.G. Ardenghi, S. Mossini

*Euonymus japonicus* Thunb. (Celastraceae)

+ (CAS) **ABR**: Pescara (Pescara), aiuola lungo Viale T. Patini (WGS84: 42.452488°N; 14.241713°E), epifita su stipite di *Phoenix canariensis*, ca. 4 m, 29 October 2017, N. Olivieri (FI). – Casual alien species new for the flora of Abruzzo.

A young individual of this species grows as an epiphyte among the stumps of the cut leafy rachis of a *Phoenix canariensis* H.Wildpret, at about 1.5 m from the ground. The settlement site is located in a coastal, partially shaded, suburban area, not far from the Adriatic Sea. In the surrounding gardens, *Euonymus japonicus* is cultivated as ornamental and the recorded plant may have originated from seeds dispersed by ornithochory.

N. Olivieri

*Freesia alba* (G.L.Mey.) Gumbl. (Iridaceae)

+ (CAS) **ABR**: San Vito Chietino (Chieti), fraz. Marina di San Vito (WGS84: 42.307275°N; 14.446527°E), prato presso giardino privato, ca. 20 m, 29 October 2017, N. Olivieri (FI). – Casual alien species new for the flora of Abruzzo.

A young individual of the species has developed in a meadow near a private garden where the species is cultivated. It grows on sandy-pelitic soil, dry in summer. The locality is close to the Adriatic Sea, and has a Mediterranean climate, but it is exposed to damp atmospheric currents coming from the sea. *Freesia alba* and its horticultural hybrids with *F. corymbosa* N.E.Br. and *F. leichtlinii* Klatt, erroneously attributed to *F. refracta* (Jacq.) Klatt (Goldblatt and Manning 2008, Galasso et al. 2018), are bulbous plants of South African origin cultivated as ornamentals, even outdoors and in the ground along the Abruzzo coasts.

N. Olivieri

*Hylotelephium spectabile* (Bureau) H.Ohba (Crassulaceae)

+ (CAS) **MAR**: Pesaro (Pesaro e Urbino), Viale della Liberazione, Mura rovereche (WGS84: 43.90999°N; 12.90483°E), parete in mattoni, con *Capparis orientalis*, *Parietaria judaica*, *Convolvulus sepium*, 28 August 2017, N. Ardenghi, S. Mossini (FI). – Casual alien species new for the flora of Marche.

Two individuals (one of which with ripening fruits) were observed on the ancient city walls of Pesaro, probably originating from the dissemination of cultivated plants in the nearby dwellings.

N.M.G. Ardenghi, S. Mossini

*Lantana camara* L. subsp. *aculeata* (L.) R.W.Sanders (Verbenaceae)

+ (CAS) **PUG**: Ugento (Lecce), fraz. Torre San Giovanni, lungo un canale artificiale presso la costa ionica (WGS84: 39.875752°N; 18.146011°E), vegetazione disturbata lungo un canale, ca. 4 m, 24 August 2017, N. Olivieri (FI). – Casual alien subspecies new for the flora of Puglia.

Some individuals grow near an artificial channel along with *Phragmites australis* (Cav.) Trin. ex Steud. subsp. *australis* in a flat coastal area, on a red soil with good water availability, characterized by disturbed vegetation and partially shaded by some *Eucalyptus camaldulensis* Dehnh. subsp. *camaldulensis*. The previous report for Puglia (Olivieri 2012) of *Lantana camara* L. for the same area of Ugento has to be referred to the same taxon recorded here. *L. camara* subsp. *aculeata* is widely cultivated for ornamental purposes and naturalized in the tropics and subtropics; in Italy it is reported in Abruzzo, Molise, Basilicata, Calabria, Sicilia, and Sardegna (Galasso et al. 2018).

N. Olivieri

*Lantana depressa* Small (Verbenaceae)

+ (CAS) **PUG**: Taranto (Taranto), presso la stazione ferroviaria (WGS84: 40.484680°N; 17.223055°E), incolto, ca. 17 m, S, 23 August 2017, N. Olivieri (FI). – Casual alien species new for the flora of Puglia.

One individual of the species grows at the base of the steps leading to an old abandoned building near the railway embankment, not far from the railway station. It has developed inside a fissure at the base of the rise of a partially eroded concrete step, in a context of ruderal vegetation dominated by *Dittrichia viscosa* (L.) Greuter subsp. *viscosa*. *Lantana depressa* is native to southern Florida and has also been reported in Sicilia (Galasso et al. 2018). According to Sanders (2012), the individual belongs to *L. depressa* var. *depressa*, originally widespread along the limestone outcrop of the Miami Rock Ridge.

N. Olivieri

***Musa basjoo* Siebold & Zucc. ex *linuma* (Musaceae)**

+ (NAT) **TOS**: Montignoso (Massa-Carrara), fraz. Cinquale, sommità dell'argine di un canale in destra del Fiume Versilia, in prossimità di una villetta residenziale (WGS84: 43.986072°N; 10.161817°E), argine di canale, 2 m, 19 July 2017, L. Lastrucci, L. Lazzaro (FI). – Naturalized alien species new for the flora of Toscana.

In the site of collection, the species is present with several individuals, of different ages. The population appears to have originated by vigorous resprout from pruning residues. Indeed, the species is widely cultivated in the gardens of the residential areas near the canals surrounding Lake Porta. Several localized stands of this species have also been observed in other sites of the Massa-Carrara and Lucca provinces.

L. Lastrucci, G. Ferretti, L. Lazzaro

***Physalis peruviana* L. (Solanaceae)**

+ (CAS) **LAZ**: Bracciano (Roma), fraz. Vigna di Valle, Museo Storico dell'Aeronautica Militare, presso l'Aeroporto di Vigna di Valle, in riva al lago (WGS84: 42.085373°N; 12.218674°E), riva di lago, 157 m, 28 July 2017, S. Buono (FI, UTV). – Casual alien species new for the flora of Lazio.

*Physalis peruviana* is a herbaceous perennial species, which has been very widely introduced across the world from South America as a cultivated plant for its fruit, as a medicinal plant, and as an ornamental (CABI 2018). It is reported as a casual alien in several administrative regions, especially in northern Italy, and as naturalized in Sicilia (Galasso et al. 2018). It is classified as an invasive plant at the global level (Global Invasive Species Database 2018). Some individuals of this species grow on sandy soil near Lake Bracciano, along with *Abutilon theophrasti* Medik, *Datura wrightii* Regel (see a previous record in this contribution), *Ludwigia peploides* (Kunth) P.H.Raven subsp. *montevidensis* (Spreng.) P.H.Raven, *Pavonia hastata* Cav. (see also Galasso et al. 2017), *Portulaca oleracea* L., *Solanum nigrum* L., and other alien species. The plants regularly develop flowers and fruits.

S. Buono, S. Magrini, A. Scoppola

***Rotala ramosior* (L.) Koehne (Lythraceae)**

+ (NAT) **EMR**: Colorno (Parma), golena del Po presso la fraz. Sacca (WGS84: 44.976183°N; 10.383083°E), fanghi di lanca, 25 m, 20 August 2017, M. Adorni (FI). – Naturalized alien species new for the flora of Emilia-Romagna.

In Galasso et al. (2018), *Rotala ramosior* is reported only for Piemonte, Lombardia, where the species grows in rice fields (Banfi and Galasso 2010), and for Veneto along the River Po (Masin and Scortegagna 2012). The population from Sacca, con-

sisting of several dozens of plants growing on muddy and damp soil, was also reported in the Acta Plantarum Forum (<http://www.floraitaliae.actaplantarum.org/viewtopic.php?t=98599>).

M. Adorni

*Styphnolobium japonicum* (L.) Schott (Fabaceae)

+ (CAS) **PUG**: Foggia (Foggia), Villa Comunale - Parco Karol Wojtyla (WGS84: 41.555366°N; 15.186515°E), ca. 64 m, 19 August 2017, N. Olivier (FI). – Casual alien species new for the flora of Puglia.

Some young individuals of the species grow along the perimetral wall of the Villa Comunale, near Via Galliani, on a calcareous alluvial vertisol subjected to partial desiccation of the herbaceous vegetation in summer, partially shaded by *Styphnolobium japonicum* and *Pinus halepensis* Mill. subsp. *halepensis*.

N. Olivier

*Trachycarpus fortunei* (Hook.) H.Wendl. (Arecaceae)

+ (CAS) **MAR**: Pesaro (Pesaro e Urbino), Viale C. Battisti (WGS84: 43.91577°N; 12.91205°E), aiuola con *Acer pseudoplatanus* coltivato, 4 m, un individuo, 28 August 2017, N. Ardenghi, S. Mossini (FI). – Casual alien species new for the flora of Marche.

A single, aged individual was found in a public flowerbed, probably grown from seeds originated from cultivated plants in the surrounding public and private gardens.

N.M.G. Ardenghi, S. Mossini

*Tradescantia fluminensis* Vell. (Commelinaceae)

+ (CAS) **MAR**: Pesaro (Pesaro e Urbino), Viale della Liberazione, Mura roveresche (WGS84: 43.90999°N; 12.90483°E), parete in mattoni, con *Capparis orientalis*, *Parietaria judaica*, *Convolvulus sepium*, 28 August 2017, N. Ardenghi, S. Mossini (FI). – Casual alien species new for the flora of Marche.

A group of plants, without flowers or fruits, was found on the ancient city walls of Pesaro, probably deriving from the dissemination of cultivated plants in the surrounding dwellings.

N.M.G. Ardenghi, S. Mossini

## Nomenclature and distribution updates from other literature sources

Nomenclature, status, and distribution updates according to Béguinot (1903), Devesa (2007), Kilian et al. (2009+), Göktürk and Sümbül (2014), Liu et al. (2017), Banfi (2018), Güzel et al. (2018), and Martini and Viciani (2018), and corrections to Galasso et al. (2018) are provided in Supplementary material 1.

G. Galasso

## Acknowledgements

We gratefully acknowledge colleagues who provided distribution, nomenclatural and taxonomic advices: Alessandro Alessandrini, Enrico Banfi, Fabrizio Bartolucci, Giordano Martini, and Daniele Viciani.

## References

- Anzalone B, Iberite M, Lattanzi E (2010) La Flora vascolare del Lazio. Informatore Botanico Italiano 42(1): 187–317.
- Ardenghi NMG, Parolo G, Di Gregorio B (2011) Notula: 60. In: Barberis G, Nepi C, Peccenini S, Peruzzi L (Eds) Notulae alla flora esotica d'Italia: 4 (54–89). Informatore Botanico Italiano 43(1): 145.
- Bacchetta G, Mayoral O, Podda L (2009) Catálogo de la flora exótica de la isla de Cerdeña (Italia). Flora Montiberica 41: 35–61.
- Banfi E (2018) A survey of the *Elymus* L. sensu lato species complex (Triticeae, Poaceae) in Italy: taxa and nothotaxa, new combinations and a key. Natural History Sciences, in press.
- Banfi E, Galasso G [Eds] (2010) La flora esotica lombarda. Museo di Storia Naturale di Milano, Milano, 274 pp. [+ CD-Rom]
- Béguinot A (1903) Dipsaceæ. In: Fiori A, Béguinot A. Flora analitica d'Italia, Vol. 3(1). Tipografia del Seminario, Padova, 141–156.
- CABI (2018) *Physalis peruviana* [original text by Parker C]. In: Invasive Species Compendium. CAB International, Wallingford. <http://www.cabi.org/isc> [accessed 05.04.2018]
- Cacciato A (1955) *Linaria heterophylla* Desf. e *Cephalaria syriaca* Schrad., nuove avventizie per Roma. Nuovo Giornale Botanico Italiano, n.ser. 61(2–3) (1954): 439–440.
- Cerutti GV, Motta A (2012) Notula: 126. In: Barberis G, Nepi C, Peccenini S, Peruzzi L (Eds) Notulae alla flora esotica d'Italia: 6 (115–136). Informatore Botanico Italiano 44(1): 189.
- D'Aleo F, Bonanno F (2016) Noterella 0176: *Datura wrightii* Regel. Acta Plantarum Notes 4: 145.
- Del Guacchio E (2011) Notula: 63. In: Barberis G, Nepi C, Peccenini S, Peruzzi L (Eds) Notulae alla flora esotica d'Italia: 4 (54–89). Informatore Botanico Italiano 43(1): 146.
- Devesa JA (2007) *Cephalaria* Schrad. ex Roem. & Schult. In: Devesa JA, Gonzalo R, Herrero A (Eds) Flora Iberica, Vol. 15. Real Jardín Botánico, CSIC, Madrid, 276–286.

- Frodin DG, Heywood VH (1968) *Cytisus* L. In: Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, Webb DA (Eds) Flora Europaea, Vol. 2. Cambridge University Press, Cambridge, 86–90.
- Galasso G, Conti F, Peruzzi L, Ardenghi NMG, Banfi E, Celesti-Grapow L, Albano A, Alessandrini A, Bacchetta G, Ballelli S, Bandini Mazzanti M, Barberis G, Bernardo L, Blasi C, Bouvet D, Bovio M, Cecchi L, Del Guacchio E, Domina G, Fascetti S, Gallo L, Gubellini L, Guiggi A, Iamónico D, Iberite M, Jiménez-Mejías P, Lattanzi E, Marchetti D, Martinetto E, Masin RR, Medagli P, Passalacqua NG, Peccenini S, Pennesi R, Pierini B, Podda L, Poldini L, Prosser F, Raimondo FM, Roma-Marzio F, Rosati L, Santangelo A, Scoppola A, Scortegagna S, Selvaggi A, Selvi F, Soldano A, Stinca A, Wagensommer RP, Wilhalm T, Bartolucci F (2018) An updated checklist of the vascular flora alien to Italy. Plant Biosystems 152(3): 556–592. <https://doi.org/10.1080/11263504.2018.1441197>
- Galasso G, Domina G, Bonari G, Buono S, Chianese G, Cortesi G, Frangini G, Iamónico D, Olivieri N, Peruzzi L, Pierini B, Roma-Marzio F, Scoppola A, Soldano A, Stinca A, Tomaselli V, Veronico G, Nepi C (2017) Notulae to the Italian alien vascular flora: 4. Italian Botanist 4: 33–41. <https://doi.org/10.3897/italianbotanist.4.21666>
- Global Invasive Species Database (2018) Species profile: *Physalis peruviana*. <http://www.iucngisd.org/gisd/speciesname/Physalis+peruviana> [accessed 05.04.2018]
- Göktürk R, Sümbül H (2014) A taxonomic revision of the genus *Cephalaria* (Caprifoliaceae) in Turkey. Turkish Journal of Botany 38(5): 927–968. <https://doi.org/10.3906/bot-1310-6>
- Goldblatt P, Manning JC (2008) The Iris Family: Natural History & Classification. Timber Press, Portland.
- Güzel ME, Kilian N, Gültepe M, Kandemir A, Coşkunçelebi K (2018) Contributions to the taxonomy of *Lactuca* (Asteraceae) in Turkey. Turkish Journal of Botany 42(2): 197–207. <https://doi.org/10.3906/bot-1707-29>
- Iocchi M, Bartolucci F (2008) La flora del litorale di Pian di Spille (Lazio settentrionale). Webbia 63(1): 135–155. <https://doi.org/10.1080/00837792.2008.10670837>
- Kilian N, Hand R, Raab-Straube E von [Eds] (2009+) Cichorieae Systematics Portal. <http://cichorieae.etaxonomy.net/portal/> [accessed 05.04.2018]
- Liu Q, Liu H, Wen J (2017) Relationships between *Sorghum bicolor* (Poaceae) and its close relatives based on genomic in situ hybridization evidence. Turkish Journal of Botany 41(1): 927–968. <https://doi.org/10.3906/bot-1606-8>
- Lucchese F [Ed.] (2017) Atlante della flora alloctona del Lazio: cartografia, ecologia e biogeografia. Vol. 1: Parte generale e flora alloctona. Regione Lazio, Direzione Ambiente e Sistemi Naturali, Roma.
- Martini G, Viciani D (2018) What happened to Linnaeus's *Iris florentina*? Re-evaluation of this taxon at species level. Taxon 67(2). <https://doi.org/10.12705/672.8>
- Masin R, Scortegagna S (2012) Flora alloctona del Veneto centro-meridionale (province di Padova, Rovigo, Venezia e Vicenza – Veneto – NE Italia). Natura Vicentina 15: 5–54.
- Matthews VA (1972) *Cephalaria* Schrader ex Roemer & Schultes. In: Davis PH (Ed.) Flora of Turkey and the East Aegean Islands, Vol. 4. Edinburgh University Press, Edinburgh, 585–597.

- Olivieri N (2012) Notula: 118. In: Barberis G, Nepi C, Peccenini S, Peruzzi L (Eds) Notulae alla flora esotica d'Italia: 6 (115–136). Informatore Botanico Italiano 44(1): 188.
- Ryves TB, Clement EJ, Foster MC (1996) Alien grasses of the British Isles. Botanical Society of the British Isles, London.
- Sanders RW (2012) Taxonomy of *Lantana* sect. *Lantana* (Verbenaceae): II. Taxonomic revision. Journal of the Botanical Research Institute of Texas 6(2): 403–441.
- Stinca A, Chianese G, D'Auria G, Del Guacchio E, Fascetti S, Perrino EV, Rosati L, Salerno G, Santangelo A (2017) New alien vascular species for the flora of southern Italy. Webbia 72(2): 295–301. <https://doi.org/10.1080/00837792.2017.1349236>
- Targioni Tozzetti A (1896) Cenni storici sulla introduzione di varie piante nell'agricoltura ed orticoltura toscana. Nuova ristampa arricchita di note lasciate manoscritte dall'autore e fatte dalla r. Società toscana di Orticoltura per cura del Dott. Eugenio Baroni. Tipografia M. Ricci, Firenze.
- Verloove F (2008) *Datura wrightii* (Solanaceae), a neglected xenophyte, new to Spain. Boute-loua 4: 37–40.
- Verloove F, Galasso G, Banfi E, Ardenghi NMG (2010) Notula: 34. In: Nepi C, Peccenini S, Peruzzi L (Eds) Notulae alla flora esotica d'Italia: 2 (22–37). Informatore Botanico Italiano 42(1): 388.
- Viegi L, Vangelisti R, D'Eugenio ML, Rizzo AM, Brilli-Cattarini A (2004) Contributo alla conoscenza della flora esotica d'Italia: le specie presenti nelle Marche. Atti della Società Toscana di Scienze Naturali, Memorie, Serie B 110(2003): 97–162.

## Supplementary material I

### Supplementary data

Edited by: Gabriele Galasso

Data type: species data

Explanation note: 1. Nomenclature updates; 2. Status/Note updates; 3 Distribution updates; 4. Synonyms, misapplied or included names.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/italianbotanist.5.25910.suppl1>